



ESTABLISHED 1978

VitaTrace Nutrition Ltd

www.vitatrace.com

WATER SOLUBLE VITAMIN PREPARATIONS PRODUCT LIST

Vitydor NG



Composition

Each 1 kg contains

Vitamin A (Retinol acetate)	IU	15'000'000
Vitamin D3 (Cholecalciferol)	IU	5'000'000
Vitamin E (dl-alpha-tocopheryl acetate)	mg	20'000
Menadione Sodium Bisulphite (Vitamin K3)	mg	4'000
Vitamin B1 (Thiamine hydrochloride)	mg	2'500
Vitamin B2 (Riboflavin)	mg	3'000
Nicotinic Acid	mg	12'000
Pantothenic Acid	mg	6'500
Vitamin B6 (Pyridoxine hydrochloride)	mg	3'000
Vitamin B12 (Cyanocobalamin)	mg	25
Folic Acid	mg	550
Biotin (H)	mg	10
Vitamin C	mg	30'000
Carrier		up to 1kg

Dosage: *Large Animals:* 100g per 1000 litres of drinking water for 5-7 days.

Poultry: 100g per 200 litres of drinking water for 5-7 days.

Purpose of use: Use in large animal and poultry water as a multivitamin additive for the prevention of vitamin deficiencies and as a nutritional booster. Can also be used during periods of stress.

Vitamin E (10%) + Se Soluble



Composition

Each 1 kg contains

Vitamin E (dl-alpha-tocopheryl acetate)	mg	100'000
Selenium (Sodium selenite)	mg	100
Carrier		up to 1kg

Dosage: *Poultry:* 1g in 2 litres of drinking water for 3-5 days.

Sheep & Goats: 1g per animal for 3-5 days.

Cattle, Horses & Camels: 5g per animal for 3-5 days.

Purpose of use: Use in drinking water for the prevention of Vitamin E and Selenium deficiencies.



Vitamin E (20%) + Se Soluble

Composition	Each 1 kg contains	
Vitamin E (dl-alpha-tocopheryl acetate)	mg	200'000
Selenium (Sodium selenite)	mg	200
Carrier		up to 1kg

Dosage: 500g per 1000 litres of drinking water for 3-5 days for poultry.

Purpose of use: Use in poultry water for the prevention of Vitamin E and selenium deficiencies, at times of stress.



B-Booster Plus soluble (B+K+Choline+Aminoacids)

Composition	Each 1 kg contains	
Vitamin B1 (Thiamine hydrochloride)	mg	5'000
Vitamin B2 (Riboflavine)	mg	6'000
Vitamin B6 (Pyridoxine hydrochloride)	mg	6'000
Vitamin B12	mg	30
Nicotinic acid (PP)	mg	40'000
Pantothenic acid (B3)	mg	20'000
Folic acid	mg	2'000
Vitamin K3	mg	5'000
Biotin (H)	mg	3
Choline	mg	20'500
Methionine	mg	10'000
Lysine	mg	20'000
Carrier		up to 1kg

Purpose of use: All cases which require immediate administration of the B-group and K vitamins and during & after administration of sulfa drugs or antibiotics in large doses.

Dosage: *Broilers and layers*: 1.25 g per litre of drinking water for 4-5 days.
Calves: 3 g per 100kg live weight, in the milk replacer, for 5 days.



Vitalyt Forte

<i>Composition</i>	<i>Each 1 kg contains</i>	
Vitamin A (Retinol acetate)	IU	10'000'000
Vitamin D3 (Cholecalciferol)	IU	1'800'000
Vitamin E (dl-a-tocopheryl acetate)	mg	3'100
Vitamin K3 (Menadione sodium bisulphite 8'400mg)	mg	4'200
Vitamin B2 (Riboflavin)	mg	2'000
Vitamin B6 (Pyridoxine hydrochloride)	mg	300
Folic Acid	mg	300
Vitamin C	mg	100'000
Glycine	mg	48'000
Monopotassium phosphate	mg	33'000
Iron	mg	9'600
Copper	mg	800
Magnesium	mg	650
Citric Acid	mg	10'000
Sodium Chloride	mg	100'000
Carrier		up to 1kg

Dosage: 0,5g - 1,0g per 1 litre of drinking water for 3-5 days for poultry.

Purpose of use: Use in poultry water as vitamins and electrolytes as anti-stress during periods of stress.



Menadione Sodium Bisulfite 20% (Vitamin K3 10%)

<i>Composition</i>	<i>Each 1 kg contains</i>	
Menadione sodium bisulfite	g	200 (Vit. K3 100 g)
Carrier		up to 1kg

Purpose of use: For the prevention of vitamin K deficiencies and at times of stress.

Dosage: Thoroughly mix 1g in 1 litre of drinking water.

Duration of treatment: 3-5 days according to instructions as only source of water.



Menadione Sodium Bisulfite 40% (Vitamin K3 20%)

Composition: *Each 1kg contains*

Menadione sodium bisulphite	mg	400'000
		(contain Vit. K3 200'000 mg)
Carrier		up to 1kg

Purpose of use: For the prevention and treatment of vitamin K deficiencies and at times of stress.

Dosage: Thoroughly mix 1g in 2 litres of drinking water for poultry.

Duration of treatment: 3-5 days according to instructions as only source of water.



Ascorbic Acid (Vitamin C) 50%

Each 1kg contains 500g of Vitamin C (Ascorbic Acid)

Purpose of use: For incorporation into poultry feed or drinking water for the prevention of Vitamin C deficiencies.

Dosage: *Poultry* : 50g in 200 litres of drinking water for 3-5 days.

Ascorbic Acid (Vitamin C) 100%

Each 1kg contains 1000g of Vitamin C (Ascorbic Acid)

Purpose of use: For incorporation into poultry feed or drinking water for the prevention of Vitamin C deficiencies.

Dosage: *Poultry* : 25g in 200 litres of drinking water for 3-5 days.



Manufactured in Cyprus, EU, by:



ESTABLISHED 1978

VitaTrace Nutrition Ltd

www.vitatracer.com

VitaTrace Nutrition Ltd

Tel: +357 22426527
 Fax: +357 22498325
 Email: reception@vitatracer.com

Postal Address:
 P.O. Box 23886, CY-1687 Nicosia, Cyprus

Offices & Laboratory:
 18 Propyleon Street, Industrial Estate, 2033, Strovolos, Cyprus

Plant:
 17 Propontidos Street, Industrial Estate, 2033, Strovolos, Cyprus